

ON PLANE GEOMETRIC SPANNERS: A SURVEY AND OPEN PROBLEMS

Prosenjit Bose*

Michiel Smid*

ABSTRACT. We review results and present open problems on different variants of the problem of constructing plane geometric spanners.

1 Introduction

Given a weighted graph $G = (V, E)$ and a real number $t \geq 1$, a t -spanner of G is a spanning subgraph G' with the property that for every edge xy in G , there exists a path between x and y in G' whose weight is no more than t times the weight of the edge xy . Thus, shortest-path distances in G' approximate shortest-path distances in the underlying graph G and the parameter t represents the approximation ratio. The smallest t for which G' is a t -spanner of G is called the *spanning ratio* of the graph G' . In the literature, the terms *stretch factor* or *dilation* are also used. The term *dilation* often refers to the more general situation where G' should additionally approximate the distance between any pair of points that lie on the *edges* of G .

Spanners have been studied in many different settings. The various settings depend on the type of underlying graph G , on the way weights are assigned to edges in G , on the specific value of the spanning ratio t , and on the function used to measure the weight of a shortest path. We concentrate on the setting where the underlying graph is geometric. In this context, a geometric graph is a weighted graph whose vertex set is a set of points in \mathbb{R}^2 and whose edge set consists of line segments joining two vertices. The edges are weighted by the Euclidean distance between their endpoints. Given a geometric graph $G = (V, E)$, a t -spanner of G is a spanning subgraph G' with the property that for every edge $xy \in G$, there is a path from x to y in G' such that the sum of the weights of the edges in this path is no more than t times $d(x, y)$, where $d(x, y)$ denotes the Euclidean distance between x and y .

There is a vast literature on different methods for constructing t -spanners with various properties in this geometric setting (see [44] for a comprehensive survey of the area). Aside from trying to build a spanner that has small spanning ratio, additional properties of the spanners are desirable. Typical goals in this area include the construction of t -spanners that

*School of Computer Science, Carleton University, Ottawa, Ontario, K1S 5B6, Canada. The authors are supported by the Natural Science and Engineering Research Council of Canada. Email: {jit,michiel}@scs.carleton.ca

also have few edges, bounded degree, fault tolerance, and low weight, to name a few. Notice that some of these properties actually oppose each other. For example, a graph with few edges or bounded degree cannot have a high fault-tolerance. Therefore, one needs to balance the various properties.

Our goal in this survey is to review results related to the following problem: Given a finite set P of points in \mathbb{R}^2 , construct a plane t -spanner of the complete geometric graph with vertex set P , for some constant $t \geq 1$. We also explore the setting where the underlying graph is not the complete graph but some other type of graph such as the unit-disk graph or the visibility graph of a set of line segments. Finally, we touch on plane spanners with bounded dilation. In addition to surveying various results related to the construction of plane t -spanners, we also mention several open problems in the area and provide some proof sketches of a few results so that the reader may get a flavor of the different techniques used in this area.

In the rest of this paper, we say that a graph is a spanner if it is a t -spanner for some constant t .

2 Plane Spanners

In what follows, unless specified otherwise, we assume that the underlying graph is the complete geometric graph. We let P be a finite set of points in the plane. The following is the central question addressed in this section: Given the point set P , is it always possible to construct a plane spanner? What constraints does planarity place on the spanning ratio?

2.1 Lower Bounds

In his seminal paper, Chew [17] was the first to study this question. First, he observed that requiring planarity does indeed impose a lower bound on the spanning ratio. Consider four points placed at the four corners of a square. Every plane geometric graph embedded on those 4 points has a spanning ratio of at least $\sqrt{2}$. This was the best known lower bound on the spanning ratio until Mulzer [42] showed that every triangulation of a regular 21-gon has spanning ratio at least $\sqrt{2.005367532} \approx 1.41611$. This leads us to our first open problem:

Open Problem 1. *What is the best lower bound on the spanning ratio of plane geometric graphs? Specifically, is there a $t > 1.41611$ and a point set P , such that every triangulation of P has spanning ratio at least t ?*

2.2 Upper Bounds

Chew [17] showed that the L_1 -metric Delaunay graph is a $\sqrt{10}$ -spanner (for the Euclidean metric). The L_1 -metric Delaunay graph of P is the plane graph that is constructed in the following way. Two points $x, y \in P$ form an edge of the Delaunay graph provided there is a point $z \in \mathbb{R}^2$ such that x and y are its nearest neighbors in the L_1 metric (see [45] for a comprehensive review of Delaunay Graphs and Voronoi Diagrams). Note that the unit circle in the L_1 metric is a diamond (a unit square tilted by 45 degrees); see Figure 1. Note that planarity of this graph is only ensured provided that no 4 points lie on the boundary of a tilted square.

Subsequently, in the journal version of [17], Chew [19] improved the result by showing that one can construct a plane graph whose spanning ratio is at most 2. He proved that a Delaunay graph built using a convex distance function defined by an equilateral triangle having one vertical side, as opposed to the L_1 -metric diamond, has a spanning ratio of 2. He refers to such equilateral triangles as *tilted equilateral-triangles* (see Figure 1). The tilted equilateral-triangle Delaunay graph of a planar point set P is constructed in the following manner. Two points $x, y \in P$ share an edge provided that there exists a tilted equilateral triangle with x and y on its boundary and no points of P in its interior.




Figure 1: Tilted equilateral triangle and L_1 -metric diamond.

2.3 Delaunay Triangulations

A natural question that Chew [19] posed is whether or not the standard (i.e., Euclidean) Delaunay triangulation is a spanner. By placing the points on the boundary of a circle, Chew noticed that the spanning ratio of the Delaunay triangulation can be at least $\pi/2 - \epsilon$ for any $\epsilon > 0$; See Figure 3(a). This led him to conjecture that not only is the standard Delaunay triangulation a spanner but that its spanning ratio is strictly less than 2. The first to show that the standard Delaunay triangulation of a point set is indeed a spanner were Dobkin

et al. [23]. They showed that the spanning ratio of the Delaunay triangulation is at most $(1 + \sqrt{5})\pi/2 \approx 5.08$.

Almost all of the proofs in the literature are constructive. We give the reader a flavor of how some of these proofs proceed by highlighting one of the cases in Dobkin *et al.*'s proof. Let $DT(P)$ be the standard Delaunay triangulation of P and $VOR(P)$ the Voronoi diagram of P . It is well known that $DT(P)$ and $VOR(P)$ are duals of each other. Given two points $x, y \in P$, construct a path from x to y in $DT(P)$, in the following way. For ease of exposition, assume that x and y are on a horizontal line, unit distance apart, with x to the left of y . Let $P_{VOR(P)}(x, y) = [x = p_1, p_2, \dots, p_k = y]$ be the sequence of sites of $VOR(P)$ whose cells intersect the segment xy ordered from x to y . We call $P_{VOR(P)}(x, y)$ a *Voronoi path*. By the duality relation between Delaunay triangulations and the Voronoi diagram, this path consists of Delaunay edges. The Voronoi path is called *one-sided* provided that all of the sites lie in one closed half-plane defined by the line containing xy . See Figure 2 for an example. Denote by c_j the intersection point of the segment xy with the Voronoi edge separating the cells of p_j and p_{j+1} . Notice that by construction the circle centered at c_j with radius $c_j p_j$, denoted $C(c_j, p_j)$, has p_j and p_{j+1} on its boundary and is empty of all other points of P . Furthermore, the arc of $C(c_j, p_j)$ defined clockwise from p_j to p_{j+1} is longer than the segment $p_j p_{j+1}$. Therefore, when $P_{VOR(P)}(x, y)$ is a one-sided Voronoi path, its length is bounded by half the boundary of the union of the circles $C(c_j, p_j)$, $j = 1, \dots, k - 1$. Since the boundary of the union of these circles has length at most π , the length of $P_{VOR(P)}(x, y)$ is at most $\pi/2$. When the Voronoi path is not one-sided, its spanning ratio can be unbounded. In this case, Dobkin *et al.* showed how to construct a path with spanning ratio at most $(1 + \sqrt{5})\pi/2$. The argument in this case is slightly more involved.




Figure 2: One-sided Voronoi path from x to y shown in bold.




Figure 3: (a) Construction providing a lower bound of $\pi/2 - \epsilon$ on the spanning ratio. (b) Construction providing lower bound of 1.581. The basic construction consists of two unit semicircles separated by a fixed distance that is optimized to maximize the spanning ratio. The shortest path from p to p' in the Delaunay triangulation has spanning ratio at least 1.581.

Subsequently, Keil and Gutwin [34] showed that the spanning ratio of the Delaunay triangulation is at most $4\pi\sqrt{3}/9 \approx 2.42$, which is currently the best known upper bound on the spanning ratio of Delaunay triangulations. Their proof is inductive and also relies heavily on the empty circle property of Delaunay triangulations. This is the best known general upper bound on the spanning ratio. Recently, Cui *et al.* [20] improved the upper bound on the spanning ratio for points in convex position. They showed that the upper bound on the spanning ratio is at most 2.33 when points are in convex position.

Open Problem 2. *What is the best upper bound on the spanning ratio of the Delaunay triangulation? Can one prove that the spanning ratio of the Delaunay triangulation is strictly less than $4\pi\sqrt{3}/9$? For points in convex position, can one prove that the spanning ratio is strictly less than 2.33?*

When considering the above problem, one also needs to consider the issue of lower bounds. Chew [19] conjectured that the worst-case spanning ratio of the Delaunay triangulation is $\pi/2$ and showed that by placing points on the boundary of a circle, one can approach this bound. The fact that one-sided Voronoi paths also have a spanning ratio of at most $\pi/2$ led many to believe that $\pi/2$ was the correct bound. Surprisingly, it was shown recently that the worst-case spanning ratio of the Delaunay triangulation is actually greater than $\pi/2$ [9]. There exists a set of points in convex position for which the spanning ratio of the Delaunay triangulation is at least 1.581. See Figure 3(b). The lower bound can be slightly improved to 1.5846 if points do not need to be in convex position. Moreover, it is shown that the lower bound on the spanning ratio of the Delaunay triangulation is essentially the same for random point sets.

Open Problem 3. *What is the best lower bound on the worst-case spanning ratio of the Delaunay triangulation? Can one construct a point set such that the spanning ratio of its Delaunay triangulation is strictly greater than 1.5846? For points in convex position, can one construct a point set such that the spanning ratio is strictly greater than 1.581?*

Although many believe that the spanning ratio of the Delaunay triangulation is strictly less than 2, no one has proven nor disproven this yet. This leads us to the following open problem.

Open Problem 4. *What is the best upper bound on the spanning ratio of a plane graph? As mentioned in Section 2.2, the best that is currently known is an upper bound of 2.*

2.4 Minimum Spanning Ratio

One question that comes to mind when contemplating the above problem is how to compute, for a given point set, the plane graph with minimum spanning ratio. The complexity of the problem is unknown, however, there is strong evidence to suggest that the problem is NP-hard. Recently, Klein and Kutz [37] showed that computing, when given a point set and a real number $t > 1$, the t -spanner with the minimum number of edges is NP-hard. In fact, Cheong *et al.* [16] showed that even computing the spanning tree with minimum spanning ratio of a given point set is an NP-hard problem. However, their proof does not imply the NP-hardness of the problem in the plane setting because the spanning tree of minimum spanning ratio need not be plane. This leads to the following two open problems:

Open Problem 5. *Is computing the plane graph of minimum spanning ratio for a given point set an NP-hard problem?*

Open Problem 6. *Is computing the plane spanning tree of minimum spanning ratio for a given point set an NP-hard problem?*

2.5 α -Diamond Spanners

The empty circle property is the key property of Delaunay triangulations that is exploited to prove the upper and lower bounds on the spanning ratio. However, there does not seem to be anything particularly special about circles. One can view the empty circle property as each edge of the triangulation having an empty *buffer region* on one side of the edge. In other words, each edge of the triangulation has a fixed proportional amount of space on one of its sides that is guaranteed to be empty of points. Das and Joseph [22] formalized this notion in the following way.

Definition 1. *Given a point set P and two points $x, y \in P$, the segment xy has the α -diamond property provided that at least one of the two isosceles triangles with base xy and base angle α does not contain any point of P . Note that the apices of the isosceles triangles need not be points of P . See Figure 4.*

A plane graph has the α -diamond property when every edge of the graph has the α -diamond property for some fixed α . Moreover, a plane graph has the d -good polygon property




Figure 4: Segment xy has the α -diamond property and segment ab does not

if for every visible pair of vertices a, b on a face f , the shortest distance from a to b around the boundary of f is at most d times the Euclidean distance between a and b . Das and Joseph showed that an α -diamond plane graph with the d -good polygon property has a spanning ratio of at most $8d\pi^2/\alpha^2 \sin^2(\alpha/4)$. This was slightly improved to $8d(\pi - \alpha)^2/\alpha^2 \sin^2(\alpha/4)$ [12]. Notice that when the value of d is 1, then a plane graph with the α -diamond property must be a triangulation. Das and Joseph showed that certain special types of triangulations possess the α -diamond property for fixed values of α . The empty circle property implies that Delaunay triangulations have the α -diamond property for $\alpha = \pi/4$. Das and Joseph also proved that the minimum weight triangulation and the greedy triangulation each have the α -diamond property with $\alpha = \pi/8$. The minimum weight triangulation of a point set is defined as the triangulation whose sum of the lengths of its edges is minimum over all possible triangulations of the given point set. It was recently shown that computing such triangulations is NP-hard [43]. The greedy triangulation is one that is constructed in the following way. Starting with just the vertices, edges are added in non-decreasing order as long as planarity is not violated. Drysdale et al. [24] improved the value of α for greedy triangulations to $\arctan(1/\sqrt{5})$. Bose *et al.* [12] subsequently improved this to $\pi/6$.

Open Problem 7. *Can the spanning ratio for plane graphs with the α -diamond property and the d -good polygon property be improved? Specifically, can one show that the spanning ratio is strictly less than $8d(\pi - \alpha)^2/\alpha^2 \sin^2(\alpha/4)$?*

Open Problem 8. *Is the α -value for greedy triangulations greater than $\pi/6$?*

Open Problem 9. *Is the α -value for minimum weight triangulations greater than $\pi/8$?*

2.6 Convex Delaunay Spanners

Chew [17, 19] showed that the Delaunay graph constructed using two different convex distance functions (based on the diamond and the tilted equilateral triangle) results in a spanner. This begs the question whether or not the Delaunay graph constructed using an arbitrary convex distance function results in a spanner. Bose *et al.* [6] answered this in the affirmative by

showing that the Delaunay graph constructed using any convex distance function results in a spanner where the spanning ratio depends on the shape of the compact convex set used to define the distance function.

2.7 Proximity-Based Plane Spanners

We already noted that it is the empty region property of Delaunay graphs that makes them spanners. The above generalizations show that the type of empty region around each edge is not particularly important but simply the fact that an empty region exists. This leads to questions such as whether or not other kinds of geometric proximity graphs are spanners. A geometric graph is a *proximity graph* when some sort of an empty region property defines the edges of the graph. Many proximity graphs are non-plane or disconnected. One well-known class of proximity graphs are the β -skeletons [35]. A β -skeleton of a set P of points in the plane, denoted $\text{BSKEL}_\beta(P)$, is a proximity graph where the proximity region for two points $x, y \in P$ is a function of β (see Figure 5):

1. For $\beta = 0$, the proximity region is the line segment \overline{xy} .
2. For $0 < \beta < 1$, the proximity region is the intersection of the two disks of radius $d(x, y)/(2\beta)$ passing through both x and y .
3. For $1 \leq \beta < \infty$, the proximity region is the intersection of the two disks of radius $\beta d(x, y)/2$ centered at the points $(1 - \beta/2)x + (\beta/2)y$ and $(\beta/2)x + (1 - \beta/2)y$.
4. For $\beta = \infty$, the proximity region is the infinite strip perpendicular to the line segment \overline{xy} .

The edge xy is in the β -skeleton of P if the proximity region of xy does not contain any other points of P . Notice that different values of the parameter β give rise to different graphs. Note also that different graphs may result for the same value of β if the proximity regions are constructed with open rather than closed disks. When the proximity region is constructed with an open disk, it is referred to as the open β -skeleton, otherwise, it is referred to as the closed β -skeleton otherwise. For the range $0 \leq \beta < 1$, the β -skeleton is not necessarily plane and for the range $\beta > 2$, the β -skeleton is not necessarily connected. The range of interest for plane connected graphs is $\beta \in [1, 2]$. The extreme values in this range define some well-known families of plane connected proximity graphs. The *Relative Neighborhood Graph* (RNG), which is also the open 2-skeleton, was first defined as a geometric graph where two vertices x, y are adjacent provided that there is no third vertex z in the graph with the property that $\max(d(x, z), d(y, z)) < d(x, y)$. The *Gabriel Graph* (GG), which is the closed 1-skeleton, was first defined as a geometric graph where two vertices x, y are adjacent provided that there is no third vertex z in the graph with the property that $d(x, z)^2 + d(z, y)^2 \leq d(x, y)^2$. It is well-known that $\text{RNG}(P) \subseteq \text{GG}(P) \subseteq \text{DT}(P)$. See Jaromczyk and Toussaint [30] for a survey of various proximity graphs.




Figure 5: The proximity regions for various values of β .

Bose *et al.* [8] were the first to study the spanning properties of β -skeletons. For the range $\beta \in [1, 2]$, they showed that the spanning ratio can be at least $(1/2 - o(1))\sqrt{n}$ and is always at most $n - 1$. They also showed that Gabriel graphs have a spanning ratio of at most $4\pi\sqrt{2n - 4}/3$ and exhibited a family of point sets where the Gabriel graph has a spanning ratio of at least $2\sqrt{n}/3$ thereby showing that the bound is $\Theta(\sqrt{n})$. They proved that the spanning ratio of the RNG is at most $n - 1$ and that there exist point sets where this bound is achieved. Moreover, even for points uniformly distributed in the unit square, they showed that the spanning ratio for the Gabriel graph and all β -skeletons with $\beta \in [1, 2]$ tends to infinity in probability as $\sqrt{\log n / \log \log n}$. Wang *et al.* [47], subsequently, showed that the spanning ratio of Gabriel graphs is at most $\sqrt{n - 1}$ and that there exist point sets where this ratio is achieved. For $\beta \in [1, 2]$, they provide an upper bound on the spanning ratio that depends on β , specifically, $(n - 1)^\gamma$ where $\gamma = 1 - \log_2(\sqrt{2/\beta})$.

Open Problem 10. For the range $\beta \in [1, 2]$, is there a stronger lower bound that depends on β ?

Open Problem 11. For the range $\beta \in [1, 2]$, is the upper bound on the spanning ratio of $(n - 1)^\gamma$, where $\gamma = 1 - \log_2(\sqrt{2/\beta})$, tight? It is tight for the GG, where $\beta = 1$ and RNG, where $\beta = 2$.

In the range $\beta \in [0, 1]$, the resulting β -skeletons are not necessarily plane. For example, the 0-skeleton is the complete graph. However, recently, Bose *et al.* [5] studied the spanning properties of the family of graphs $\text{RGG}_\beta(P) := \text{BSKEL}_\beta(P) \cap \text{DT}(P)$ for $\beta \in [0, 1]$ called *Relaxed Gabriel Graphs*. Since $\text{GG}(P) \subseteq \text{BSKEL}_\beta(P) \cap \text{DT}(P)$, this family of graphs is connected and plane, hence the name *Relaxed Gabriel graphs* for this family. They showed upper and lower bounds on the spanning ratio for $\text{BSKEL}_\beta(P)$ that depend on $\beta \in [0, 1]$. The upper bound on the spanning ratio of $\text{RGG}_\beta(P)$ is $O(n^\gamma)$, where $\gamma = \frac{1}{2} - \frac{1}{2} (\log(1 + \cos(\frac{\alpha}{2})))$ and $\beta = \sin(\alpha/2)$. For the lower bound, they showed that there exist point sets where the spanning ratio of $\text{RGG}_\beta(P)$ is $\Omega(n^\gamma)$, with $\gamma = \frac{1}{2} - \frac{1}{2} \left(\log \left(1 + \sqrt{\frac{1 + \cos(\frac{\alpha}{2})}{2}} \right) \right)$ and $\beta = \sin(\alpha/2)$.

Open Problem 12. Are these upper and lower bounds tight?

2.8 Bounded-Degree Plane Spanners

Another question that comes to mind is whether or not it is possible to build a plane spanner with bounded degree. All of the above plane spanners can have unbounded degree.

Bose *et al.* [10] were the first to show the existence of a plane t -spanner (for some constant t), whose maximum vertex degree is bounded by a constant. To be more precise, they showed that the Delaunay triangulation of any set P of points in the plane contains a subgraph, which is a t -spanner for P , where $t = 4\pi(\pi + 1)\sqrt{3}/9$, and whose maximum degree is at most 27. Subsequently, Li and Wang [40] reduced the degree bound to 23 by showing the following: For any real number γ with $0 < \gamma \leq \pi/2$, the Delaunay triangulation contains a subgraph that is a t -spanner, where $t = \max\{\frac{\pi}{2}, 1 + \pi \sin \frac{\gamma}{2}\} \cdot \frac{4\pi\sqrt{3}}{9}$, and whose maximum degree is at most $19 + \lceil 2\pi/\gamma \rceil$. For $\gamma = \pi/2$, the degree bound is 23. In [14], Bose *et al.* improved the degree bound to 17 and generalized the result to α -diamond triangulations (i.e. they showed that every α -diamond triangulation contains a subgraph that is a plane bounded-degree spanner). Currently, the degree bound stands at 14 as shown by Kanj and Perkovic [31]; again, their spanner is a subgraph of the Delaunay triangulation.

The obvious open questions in this area are:

Open Problem 13. *What is the smallest maximum degree that can be achieved for plane spanners that are subgraphs of the Delaunay triangulation?*

Open Problem 14. *What is the smallest maximum degree that can be achieved for plane spanners?*

If one does not insist on having a plane spanner, then it is possible to construct a spanner with maximum degree 3 [21]. It is easy to see that there exist point sets such that every graph of maximum degree 2 defined on that point set has unbounded spanning ratio. As such, the more interesting question becomes whether the planarity constraint actually imposes a higher lower bound on the maximum degree. Thus, we have the following open problem:

Open Problem 15. *With the constraint of constructing a plane spanner, is there a lower bound on the maximum degree that is greater than 3? That is, can we show the following: For every real number $t > 1$, there exists a set P of points, such that every plane degree-3 spanning graph of P has spanning ratio greater than t .*

3 Low Weight Plane Spanners

The weight of a graph is the sum of the weights of its edges. A lower bound on the weight of a connected graph is the weight of the minimum spanning tree. A graph is said to have *low weight* if its weight is $O(1)$ times the weight of the minimum spanning tree. As such, one objective in this area is to build plane spanners that have constant spanning ratio and

low weight. Gudmundsson *et al.* [29] presented a very general method to compute, for any given spanner G , a subgraph of G whose spanning ratio is at most $1 + \epsilon$ times that of G and whose weight is at most $O(1)$ times the weight of the minimum spanning tree. The constant hidden in the $O(1)$ is fairly large. When considering plane graphs specifically, Levkopoulos and Lingas [38] showed that, for any given real number $r > 2$, the Delaunay triangulation can be used to construct a plane graph that is a t -spanner for $t = (r - 1)4\pi\sqrt{3}/9$ and whose total weight is at most $1 + 2/(r - 2)$ times the weight of a minimum spanning tree. Subsequently, Kanj *et al.* [32] showed how this method can be generalized to build bounded degree plane spanners. They showed that for any integer constant $k \geq 14$ and $r > 2$, one can build a plane t -spanner with maximum degree k , where $t = (r - 1)(4\pi\sqrt{3}/9)/(1 + 2\pi(k \cos(\pi/k)))$ and whose total weight is at most $1 + 2/(r - 2)$ times the weight of the minimum spanning tree.

Open Problem 16. *Are these bounds tight?*

4 $(1 + \epsilon)$ -Plane Steiner Spanners

As we have seen in Section 2.1, there do not necessarily always exist plane graphs that have a spanning ratio arbitrarily close to 1. Let P be a set of n points in the plane. In this section, we consider plane Steiner spanners, which are plane graphs whose vertex sets contain P . Vertices that do not belong to P are called Steiner points. It turns out that by allowing $O(n)$ Steiner points, we can obtain a spanning ratio of $1 + \epsilon$, for any fixed $\epsilon > 0$. We emphasize that the spanning ratio is defined only in terms of point-pairs in the set P .

We give a sketch of the construction, which is due to Arikati *et al.* [2]. The starting point is to consider the plane Steiner spanner problem in the L_1 metric. Thus, we want to construct a plane graph, whose vertex set contains P , such that any two points x and y of P are connected by a path whose L_1 length is at most $1 + \epsilon$ times the L_1 distance between x and y .

A box is defined to be an axes-parallel square whose longest side is at most twice as long as its shortest side. A doughnut is defined to be the set-theoretic difference $B \setminus B'$ of two boxes B and B' , where B' is contained in B ; see Figure 6. We put the following additional restriction on any doughnut: For any edge e' of the inner box B' , consider the corresponding edge e of the outer box B ; for example, if e' is the top edge of B' , then e is the top edge of B . We require that the distance between e and e' is either zero or at least the length of e' .

Let C be a square that contains all points of P . Using an algorithm of Arya *et al.* [3], we can compute, in $O(n \log n)$ time, a subdivision of C into $O(n)$ cells, such that each cell is either a box or a doughnut. Moreover, each box contains at most one point of P , whereas each doughnut is empty of points. Notice that the edges of this subdivision define a graph. We augment this graph in the following way.

Consider a box B in the subdivision and let ℓ be the length of its shortest side. On each




Figure 6: The doughnut $B \setminus B'$.

side of B , we place $O(1/\epsilon)$ Steiner points such that successive Steiner points have distance $\epsilon\ell$. Then we connect these Steiner points by a grid; see Figure 7. If B contains a point, say p , of P , then we also add horizontal and vertical rays from p to the edges of B . By turning all intersection points into Steiner points, we obtain a plane graph inside B consisting of $O(1/\epsilon^2)$ vertices. For each doughnut $B \setminus B'$, we do a similar construction, one for the outer box B and one for the inner box B' , as illustrated in Figure 7. Again, this results in a plane graph inside $B \setminus B'$ consisting of $O(1/\epsilon^2)$ vertices.

Since the number of cells in the subdivision is $O(n)$, we obtain a plane graph whose vertex set contains P and that has $O(n/\epsilon^2)$ Steiner points. Let x and y be two points of P , and consider the Manhattan path M (with two segments) between x and y . It is not difficult to see that our graph contains a path between x and y whose length is at most $1 + O(\epsilon)$ times the length of M . Thus, our graph is a plane Steiner $(1 + O(\epsilon))$ -spanner of the point set P .

To obtain a plane Steiner $(1 + \epsilon)$ -spanner for the Euclidean metric, we take $O(1/\epsilon)$ coordinate systems, obtained by rotating the X - and Y -axes by angles of $i\epsilon$, for $0 \leq i < 2\pi/\epsilon$. For each coordinate system, we construct the plane Steiner spanner for the L_1 metric corresponding to that system. By overlaying all these graphs, Arikati et al. [2] show that we obtain a plane graph with $O(n/\epsilon^4)$ Steiner points. Since for any two points x and y in P , there




Figure 8: All triangulations with spanning ratio 1.

six vertices. Eppstein [27] has shown that these are the only triangulations with spanning ratio 1. Notice that any finite point set P that is contained in the vertex set of one of these triangulations has dilation 1. Klein and Kutz [36] have shown that the dilation of every other point set is strictly larger than 1.

Ebbers-Baumann et al. [26] have shown that if P is the (infinite) set of points on a closed convex curve, then its dilation is larger than 1.00157. They have also shown that the dilation of every finite point set is less than 1.1247.

Open Problem 18. *What is the smallest value of t such that every finite set of points in the plane has dilation at most t ? It is known that $1.00157 \leq t \leq 1.1247$.*

It turns out that even for small point sets, it is very difficult to determine their dilation:

Open Problem 19. *What is the dilation of the vertices of a regular 5-gon? It is known that this dilation is at most 1.02046; see [26].*

We now turn to the so-called geometric dilation. Consider a plane graph and let x and y be two distinct points “on” G , i.e., each of x and y is either a vertex or in the interior of some edge. Let $SP_G(x, y)$ be the ratio of the shortest-path distance between x and y in G to the Euclidean distance $d(x, y)$. The geometric dilation of G is defined to be the supremum of $SP_G(x, y)$ over all such pairs x, y . For example, consider a triangle T and let α be its largest acute angle. If we consider T to be a graph whose vertex set consists of the three vertices of T , then the spanning ratio of T is equal to 1. On the other hand, the geometric dilation of T is equal to $\sqrt{2/(1 - \cos \alpha)}$, which is at least 2.

The geometric dilation of a finite set P of points in the plane is defined to be the infimum of the geometric dilations of all finite plane graphs whose vertex sets contain P .

As opposed to the dilation of the vertex set of the regular n -gon P_n , its geometric dilation has been determined by Ebbers-Baumann et al. [25]: The geometric dilations of P_3 and P_4 are $2/\sqrt{3}$ and $\sqrt{2}$, respectively. For $n \geq 5$, the geometric dilation of P_n is equal to $\pi/2$.

In [25], it is also shown that the geometric dilation of every finite point set is less than 1.678. Furthermore, they showed that the geometric dilation of the vertices of the 19×19 grid is larger than $(1 + 10^{-11})\pi/2$.

Open Problem 20. *What is the smallest value of t such that every finite set of points in the plane has geometric dilation at most t ? It is known that $(1 + 10^{-11})\pi/2 \leq t \leq 1.678$.*

6 Variants

In the previous sections, the underlying graph has been the complete geometric graph. We now review some results when the underlying graph is not necessarily the complete geometric graph but a subgraph. We explore two different settings: one where the underlying graph is the visibility graph of a set of line segments and the other where the underlying graph is the unit-disk graph. We begin with the former.

6.1 Constrained Setting

Before we can review the results in the constrained setting, we need to outline precisely what is meant by the constrained setting. Let P be a set of points in the plane and let L be a set of non-crossing line segments whose endpoints are in P . Two points x and y of P are *visible* with respect to L provided the segment xy does not properly intersect any segment of L . Two line segments intersect properly if they share a common interior point. The visibility graph of P constrained to L , denoted $\text{VIS}(P, L)$, is the geometric graph whose vertex set is P and whose edge set contains L as well as one edge for each visible pair of vertices (See Figure 9). All edges, including the constrained edges, are weighted by their length. A spanning subgraph of $\text{VIS}(P, L)$ whose edge set contains L is a geometric graph *constrained to L* . In such a graph, the elements of L are referred to as the *constrained edges*, whereas all other edges are referred to as *unconstrained edges* or *visibility edges*. The underlying graph in this subsection is $\text{VIS}(P, L)$. Thus, a constrained geometric graph $G(P, L)$ is a *constrained t -spanner* of $\text{VIS}(P, L)$ provided that for every edge xy in $\text{VIS}(P, L)$, the length of the shortest path between x and y in $G(P, L)$ is at most t times the Euclidean distance between x and y . Note that if $G(P, L)$ is a constrained t -spanner, then for every pair x, y of points in P (not just visible edges), the shortest path from x to y in $G(P, L)$ is at most t times the shortest path from x to y in $\text{VIS}(P, L)$. The fundamental question to address here is: Given $\text{VIS}(P, L)$, does there always exist a plane constrained spanner $G(P, L)$ of $\text{VIS}(P, L)$?

Chew [17, 19] mentioned that his technique (see in Section 2.2) can be extended to




Figure 9: The visibility graph $\text{Vis}(P, L)$ where segments of L are shown in bold.

the constrained setting. Karavelas [33] noted that the proof in Dobkin *et al.* [23] can also be extended to the constrained setting, thereby asserting that the *Constrained Delaunay triangulation*, denoted $\text{CDT}(P, L)$, is a $(1 + \sqrt{5})\pi/2$ -spanner of $\text{Vis}(P, L)$. The constrained Delaunay triangulation was independently introduced by Chew [18] and Wang and Schubert [46]. It is a generalization of the standard Delaunay triangulation. Two points $x, y \in P$ form an edge in $\text{CDT}(P, L)$ provided that there exists a disk with x and y on its boundary that does not contain any point $z \in P$ that is visible to both x and y . Subsequently, Bose and Keil [11] proved that the spanning ratio of the $\text{CDT}(P, L)$ is most $4\pi\sqrt{3}/9$. Recently, Bose *et al.* [12] generalized the results of Das and Joseph [22] to the constrained setting. A constrained graph $G(P, L)$ is said to have the *visible α -diamond property* if, for every unconstrained edge e in the graph, at least one of the two isosceles triangles, with e as the base and base angle α , does not contain any points of P visible to the endpoints of e . Refer to Figure 10. Furthermore, $G(P, L)$ has the *d -good polygon property* if for every visible pair of vertices a and b on a face f , the shortest distance from a to b around the boundary of f is at most d times the Euclidean distance between a and b . Bose *et al.* [12] generalized the results on α -diamond spanners in the following way: Given fixed $\alpha \in (0, \pi/2)$ and $d \geq 1$, if a constrained plane graph $G(P, L)$ has both the visible α -diamond property and the d -good polygon property, then its spanning ratio is at most $\frac{8(\pi-\alpha)^2 d}{\alpha^2 \sin^2(\alpha/4)}$.




Figure 10: The edge e has the visible α -diamond property.

Open Problem 21. *Essentially, all of the questions that are open in the unconstrained setting are also open in the constrained setting since the constrained setting is a generalization of the*

unconstrained one.

6.2 Unit-Disk Graphs

Given a set P of points in the plane, the unit-disk graph, denoted $\text{UDG}(P)$, is the geometric graph whose vertex set is P with two vertices being joined by an edge provided the length of the edge is at most a specified unit. There has been much interest in studying spanners of $\text{UDG}(P)$ in the wireless network community since these graphs are often used to model wireless adhoc networks (see [4] for an overview of the area).

The question to address in this area is: Does there always exist a plane spanner of the unit-disk graph? Before answering this question, we first highlight a connection between *strong spanners* and spanners of the unit-disk graph. Given a geometric graph G , a t -spanner G' of G is *strong* if for every edge $xy \in G$, there is a path from x to y in G' such that the sum of the distances of the edges in this path is no more than t times $d(x, y)$ and every edge on this path has length at most $d(x, y)$. It is the latter property that distinguishes a spanner from a strong spanner. Note that any method for constructing a plane strong spanner of the complete geometric graph becomes a technique for constructing a plane spanner of the unit-disk graph. That is, if a graph G is a plane strong spanner of the complete geometric graph, then $G \cap \text{UDG}$ is a plane strong spanner of UDG , provided UDG is connected.

Bose *et al.* [13] were the first to show that one can construct a strong plane spanner of the complete geometric graph by proving that the Delaunay triangulation is a strong $4\pi\sqrt{3}/9$ -spanner. They showed that the proof by Keil and Gutwin [34] can be generalized to show that the Delaunay triangulation is indeed a strong spanner. This implies that $\text{DT}(P) \cap \text{UDG}(P)$ is a plane spanner of $\text{UDG}(P)$. In [14], Bose *et al.* showed that one can construct a bounded-degree plane spanner of $\text{UDG}(P)$ where the maximum degree is at most 17.

The algorithms to obtain the above results are all *centralized*. This means that the algorithm is aware of the whole graph. Since there is interest in studying these spanners in the wireless network community, it should be noted that centralized algorithms are undesirable in that setting. In the wireless network setting, the challenge is to compute these spanners in a *local* manner. A wireless ad hoc network consists of a set P of n wireless nodes in the plane. Each wireless node $u \in P$ can only communicate directly with nodes that are within its communication range. If we assume that this range is equal to one unit for each node, then one can see how the unit-disk graph $\text{UDG}(P)$ models a wireless ad hoc network. We now describe what it means for an algorithm to be *local* in this setting.

In the wireless setting, it is assumed that $\text{UDG}(P)$ is connected and that each node knows its position and the position of all its neighbors within its communication range. The position information (or some other identifier) is used to distinguish the nodes. For simplicity, a message is often defined as the x -coordinate and y -coordinate of a point since that is the content of most messages in the algorithms described below. Nodes communicate with each

other by broadcasting messages and the metric used to measure the performance of different construction algorithms is the total number of messages broadcast as a function of the number of nodes. The construction algorithm is usually synchronized and a communication round is defined as the period between the sending of a message and the complete processing of the message on the receiver side. For any positive integer k , let $N_k(v) = \{w \mid \text{there is a path in } \text{UDG}(P) \text{ between } v \text{ and } w \text{ with at most } k \text{ edges}\}$. If every node v can compute the value of any computable function with domain $N_k(v)$ by an algorithm, we define this algorithm to be a k -local algorithm. For example, let $DT(P)$ be a function that returns the Delaunay triangulation of P , then the localized algorithm where each node v runs $DT(N_2(v))$ is a 2-local algorithm. Computing $N_k(v)$ is typically more expensive than computing $N_{k-1}(v)$ in a localized environment, therefore, it is desirable to design k -local algorithms with the smallest k as possible.

Let $\text{UDEL}(P)$ be the intersection between the unit-disk graph and the Delaunay triangulation of P . Gao *et al.* [28] proposed a localized algorithm to build a plane graph called the restricted Delaunay graph (RDG), which is a supergraph of $\text{UDEL}(P)$. In RDG, each node u maintains a set $E(u)$ of edges incident on u . These edges in $E(u)$ satisfy that 1) each edge in $E(u)$ is at most one unit; 2) the edges are consistent, i.e., $uv \in E(u)$ if and only if $uv \in E(v)$; 3) the graph obtained is plane; and 4) all the Delaunay edges with length at most one are guaranteed to be in the union of the $E(u)$'s. However, the total message complexity of their algorithm is $O(n^2)$.

In [39], Li *et al.* defined a k -localized Delaunay triangle as a triangle Δuvw whose interior of the circumcircle $\text{disk}(u, v, w)$ does not contain any node of $N_k(u)$, $N_k(v)$ or $N_k(w)$, and all edges of the triangle Δuvw have a length of no more than one unit. They also defined the k -localized Delaunay graph, denoted by $\text{LDel}^{(k)}(P)$, as the graph that contains exactly all Gabriel edges with length at most one and the edges of all k -localized Delaunay triangles. They showed that $\text{LDel}^{(k)}(P)$ is a supergraph of $\text{UDEL}(P)$ and plane if $k \geq 2$. They also defined $\text{PLDel}(P)$ as the plane graph obtained by removing intersecting edges, which do not belong to $\text{LDel}^{(2)}(P)$, from $\text{LDel}^{(1)}(P)$. Notice that $\text{PLDel}(P)$ is a subgraph of $\text{LDel}^{(1)}(P)$, a supergraph of $\text{LDel}^{(2)}(P)$ and also a t -spanner of $\text{UDG}(P)$ for $t = \frac{4\pi\sqrt{3}}{9}$.

In [39], Li *et al.* also proposed a 3-local algorithm to compute $\text{PLDel}(P)$ with total communication cost of $O(n)$. If we assume that sending each ID costs at most one message, the constant in the $O(n)$ bound is at most 49. Also, their algorithm needs four communication rounds. Araújo *et al.* [1] improved the work of Li *et al.* [39] by proposing a fast 2-local algorithm to compute $\text{PLDel}(P)$. Their algorithm only needs one communication round and the message complexity can be bounded by $11n$. Bose *et al.* [7] improved the work of Araújo *et al.* [1] by presenting an efficient 2-local algorithm to compute $\text{PLDel}(P)$. Their algorithm needs one communication round and the message complexity can be bounded by $5n$.

Open Problem 22. *Can one reduce the total number of messages in order to compute a plane local spanner of $\text{UDG}(P)$?*

Wang and Li [48] presented a 3-local algorithm that computes a plane spanner of

UDG(P) with degree bound 23. The communication cost of their algorithm is $O(n)$. Very recently, Kanj *et al.* [32] presented, for any given $k \geq 14$ and $\lambda > 2$, a $\lfloor (8/\pi)(\lambda + 1)^2 \rfloor$ -local algorithm that constructs a plane spanner of UDG(P) with degree bound k . Both of these two algorithms are based on the construction of the 2-localized Delaunay graph $LDel^{(2)}(P)$. Although $LDel^{(2)}(P)$ could be constructed with communication cost $O(n)$ [48] by using the result of [15], the constant in the $O(n)$ bound could be several hundred.

References

- [1] F. Araújo and L. Rodrigues. Fast localized Delaunay triangulation. In *Proceedings of the 8th International Conference on Principles of Distributed Systems (OPODIS 2004)*, volume 3544 of *Lecture Notes in Computer Science*, pages 81–93, Berlin, 2005. Springer-Verlag.
- [2] S. Arikati, D. Z. Chen, L. P. Chew, G. Das, M. Smid, and C. D. Zaroliagis. Planar spanners and approximate shortest path queries among obstacles in the plane. In *Proceedings of the 4th European Symposium on Algorithms*, volume 1136 of *Lecture Notes in Computer Science*, pages 514–528, Berlin, 1996. Springer-Verlag.
- [3] S. Arya, D. M. Mount, N. S. Netanyahu, R. Silverman, and A. Wu. An optimal algorithm for approximate nearest neighbor searching in fixed dimensions. *Journal of the ACM*, 45:891–923, 1998.
- [4] M. Barbeau and E. Kranakis. *Principles of Ad Hoc Networking*. Wiley, 2007.
- [5] P. Bose, J. Cardinal, S. Collette, E. Demaine, B. Palop, P. Taslakian, and N. Zeh. Relaxed Gabriel graphs. In *Proc. Canadian Conf. on Computational Geometry*, 2009, to appear.
- [6] P. Bose, P. Carmi, S. Collette, and M. Smid. On the stretch factor of convex Delaunay graphs. In *ISAAC*, pages 656–667, 2008.
- [7] P. Bose, P. Carmi, M. Smid, and D. Xu. Communication-efficient construction of the plane localized Delaunay graph. *CoRR*, abs/0809.2956, 2008.
- [8] P. Bose, L. Devroye, W. Evans, and D. Kirkpatrick. On the spanning ratio of Gabriel graphs and β -skeleton. *SIAM Journal on Discrete Mathematics*, 20(2):412–427, 2006.
- [9] P. Bose, L. Devroye, M. Löffler, J. Snoeyink, and V. Verma. The spanning ratio of the Delaunay triangulation is greater than $\pi/2$. In *Proc. Canadian Conf. on Computational Geometry*, 2009, to appear.
- [10] P. Bose, J. Gudmundsson, and M. Smid. Constructing plane spanners of bounded degree and low weight. *Algorithmica*, 42:249–264, 2005. Preliminary version appeared in *ESA* 2002.
- [11] P. Bose and M. Keil. On the stretch factor of the constrained Delaunay triangulation. In *Proc. of the Int. Symp. on Voronoi Diagrams in Science and Engineering (ISVD)*, pages 25–31, 2006.

- [12] P. Bose, A. Lee, and M. Smid. On generalized diamond spanners. In *Proc. Workshop on Algorithms and Data Structures*, pages 325–336, 2007.
- [13] P. Bose, A. Maheshwari, G. Narasimhan, M. Smid, and N. Zeh. Approximating geometric bottleneck shortest paths. *Computational Geometry: Theory and Applications*, 29(3):233–249, 2004.
- [14] P. Bose, M. Smid, and D. Xu. Delaunay and diamond triangulations contain spanners of bounded degree. *International Journal of Computational Geometry and Applications*, 19:119–140, 2009.
- [15] G. Calinescu. Computing 2-hop neighborhoods in ad hoc wireless networks. In *Proceedings of ADHOC-NOW'03*, volume 2865 of *Lecture Notes in Computer Science*, pages 175–186, Berlin, 2003. Springer-Verlag.
- [16] O. Cheong, H. Haverkort, and M. Lee. Computing a minimum-dilation spanning tree is NP-hard. *Computational Geometry: Theory and Applications*, 41:188–205, 2008.
- [17] P. Chew. There is a planar graph almost as good as the complete graph. In *Proc. ACM Symposium on Computational Geometry*, pages 169–177, 1986.
- [18] P. Chew. Constrained Delaunay triangulations. In *Proc. 3rd Annu. ACM Sympos. Comput. Geom.*, pages 215–222, 1987.
- [19] P. Chew. There are planar graphs almost as good as the complete graph. *Journal of Computer and System Sciences*, 39:205–219, 1989.
- [20] S. Cui, I. Kanj, and G. Xia. On the dilation of Delaunay triangulations of points in convex position. In *Proc. Canadian Conf. on Computational Geometry*, 2009, to appear.
- [21] G. Das and P. J. Heffernan. Constructing degree-3 spanners with other sparseness properties. *Int. J. Found. Comput. Sci.*, 7(2):121–136, 1996.
- [22] G. Das and D. Joseph. Which triangulations approximate the complete graph? In *Proc. International Symposium on Optimal Algorithms*, volume 401 of *Lecture Notes in Computer Science*, pages 168–192. Springer-Verlag, 1989.
- [23] D. P. Dobkin, S. J. Friedman, and K. J. Supowit. Delaunay graphs are almost as good as complete graphs. *Discrete Computational Geometry*, 5:399–407, 1990.
- [24] R. Drysdale, G. Rote, and O. Aichholzer. *A simple linear time greedy triangulation algorithm for uniformly distributed points*. Report IIG-408, Institutes for Information Processing, Technische Universität Graz, 1995.
- [25] A. Ebbers-Baumann, A. Grüne, and R. Klein. The geometric dilation of finite point sets. *Algorithmica*, 44:137–149, 2006.
- [26] A. Ebbers-Baumann, A. Grüne, R. Klein, M. Karpinski, C. Knauer, and A. Lingas. Embedding point sets into plane graphs of small dilation. *International Journal of Computational Geometry & Applications*, 17:201–230, 2007.

- [27] D. Eppstein. The Geometry Junkyard. <http://www.ics.uci.edu/~eppstein/junkyard/dilation-free>.
- [28] J. Gao, L. J. Guibas, J. Hershberger, L. Zhang, and A. Zhu. Geometric spanners for routing in mobile networks. *IEEE Journal on Selected Areas in Communications Special issue on Wireless Ad Hoc Networks*, 23(1):174–185, 2005.
- [29] J. Gudmundsson, C. Levcopoulos, and G. Narasimhan. Fast greedy algorithms for constructing sparse geometric spanners. *SIAM J. Comput.*, 31(5):1479–1500, 2002.
- [30] J. Jaromczyk and G. Toussaint. Relative neighborhood graphs and their relatives. *Proceedings of the IEEE*, 80(9):1502–1517, 1992.
- [31] I. A. Kanj and L. Perkovic. On geometric spanners of Euclidean and unit disk graphs. In *Symposium on Theoretical Aspects of Computer Science*, pages 409–420, 2008.
- [32] I. A. Kanj, L. Perkovic, and G. Xia. Computing lightweight spanners locally. In *Proceedings of Distributed Computing (DISC)*, pages 365–378, 2008.
- [33] M. I. Karavelas. *Proximity Structures for Moving Objects in Constrained and Unconstrained Environments*. Ph.D. thesis, Dept. Comput. Sci., Stanford University, 2001.
- [34] J. M. Keil and C. A. Gutwin. Classes of graphs which approximate the complete Euclidean graph. *Discrete Computational Geometry*, pages 13–28, 1992.
- [35] D. Kirkpatrick and J. Radke. A framework for computational morphology. In *Computational Geometry*, pages 217–248, 1985.
- [36] R. Klein and M. Kutz. The density of iterated crossing points and a gap result for triangulations of finite point sets. In *Proceedings of the 22nd ACM Symposium on Computational Geometry*, pages 264–272, 2006.
- [37] R. Klein and M. Kutz. Computing geometric minimum-dilation graphs is NP-hard. In *Proceedings of the 14th International Symposium on Graph Drawing (GD 2006)*, volume 4372 of *Lecture Notes in Computer Science*, pages 196–207, Berlin, 2007. Springer-Verlag.
- [38] C. Levcopoulos and A. Lingas. There are planar graphs almost as good as the complete graphs and almost as cheap as minimum spanning trees. *Algorithmica*, 8:251–256, 1992.
- [39] X.-Y. Li, G. Calinescu, P.-J. Wan, and Y. Wang. Localized Delaunay triangulation with applications in wireless ad hoc networks. *IEEE Transactions on Parallel and Distributed Systems*, 14(10):1035–1047, 2003.
- [40] X.-Y. Li and Y. Wang. Efficient construction of low weighted bounded degree planar spanner. *International Journal of Computational Geometry & Applications*, 14:69–84, 2004.
- [41] A. Maheshwari, M. Smid, and N. Zeh. I/O-efficient algorithms for computing planar geometric spanners. *Computational Geometry: Theory and Applications*, 40:252–271, 2008.

- [42] W. Mulzer. *Minimum Dilation Triangulations for the Regular n -gon*. Masters Thesis, Freie Universität Berlin, 2004.
- [43] W. Mulzer and G. Rote. Minimum-weight triangulation is NP-hard. *J. ACM*, 55(2), 2008.
- [44] G. Narasimhan and M. Smid. *Geometric Spanner Networks*. Cambridge University Press, 2007.
- [45] A. Okabe, B. Boots, K. Sugihara, and S. Chiu. *Spatial Tessellations: Concepts and Applications of Voronoi Diagrams, 2nd edition*. John Wiley and sons, 2000.
- [46] C. A. Wang and L. Schubert. An optimal algorithm for constructing the Delaunay triangulation of a set of line segments. In *Proc. 3rd Annu. ACM Sympos. Comput. Geom.*, pages 223–232, 1987.
- [47] W. Wang, X.-Y. Li, Y. Wang, and W.-Z. Song. The spanning ratios of β -skeleton. In *cccg03*, pages 35–38, 2003.
- [48] Y. Wang and X.-Y. Li. Localized construction of bounded degree and planar spanner for wireless ad hoc networks. *Mobile Networks and Applications*, 11(2):161–175, 2006.